



Reaction Ensemble Molecular Dynamics: Direct Simulation of the Dynamic Equilibrium Properties of Chemically Reacting Mixtures

by John K. Brennan, Martin Lísal, Keith E. Gubbins, and Betsy M. Rice

ARL-RP-144

September 2006

A reprint from *Physical Review E*, vol. 70, pp. 061103-1–061103-4, 2004.

Reprinted with permission from *Physical Review E*. Copyright 2004,
the American Physical Society.

NOTICES

Disclaimers

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

Citation of manufacturer's or trade names does not constitute an official endorsement or approval of the use thereof.

Destroy this report when it is no longer needed. Do not return it to the originator.

Army Research Laboratory

Aberdeen Proving Ground, MD 21005-5066

ARL-RP-144**September 2006**

Reaction Ensemble Molecular Dynamics: Direct Simulation of the Dynamic Equilibrium Properties of Chemically Reacting Mixtures

John K. Brennan and Betsy M. Rice
Weapons and Materials Research Directorate, ARL

Martin Lísal
Institute of Chemical Process Fundamentals and
J. E. Purkyně University, Czech Republic

Keith E. Gubbins
North Carolina State University, NC

A reprint from *Physical Review E*, vol. 70, pp. 061103-1–061103-4, 2004.

Reprinted with permission from *Physical Review E*. Copyright 2004,
the American Physical Society.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
<p>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p> <p>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</p>					
1. REPORT DATE (DD-MM-YYYY) September 2006		2. REPORT TYPE Reprint		3. DATES COVERED (From - To) February 2004–October 2004	
4. TITLE AND SUBTITLE Reaction Ensemble Molecular Dynamics: Direct Simulation of the Dynamic Equilibrium Properties of Chemically Reacting Mixtures				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) John K. Brennan, Martin Lísal,* Keith Gubbins,† and Betsy M. Rice				5d. PROJECT NUMBER H4311	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army Research Laboratory ATTN: AMSRD-ARL-WM-BD Aberdeen Proving Ground, MD 21005-5066				8. PERFORMING ORGANIZATION REPORT NUMBER ARL-RP-144	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.					
13. SUPPLEMENTARY NOTES <p>*E. Hála Laboratory of Thermodynamics, Institute of Chemical Process Fundamentals, Academy of Sciences of the Czech Republic, 165 02 Prague 6, Czech Republic, and Department of Physics, J. E. Purkyně University, 400 96 Usti n. Lab., Czech Republic</p> <p>†North Carolina State University, Department of Chemical Engineering, Raleigh, NC 27695-7905</p> <p>A reprint from <i>Physical Review E</i>, vol. 70, pp. 061103-1–061103-4, 2004.</p>					
14. ABSTRACT <p>A molecular simulation method to study the dynamics of chemically reacting mixtures is presented. The method uses a combination of stochastic and dynamic simulation steps, allowing for the simulation of both thermodynamic and transport properties. The method couples a molecular dynamics simulation cell (termed dynamic cell) to a reaction mixture simulation cell (termed control cell) that is formulated upon the reaction ensemble Monte Carlo (RxMC) method, hence the term reaction ensemble molecular dynamics. Thermodynamic and transport properties are calculated in the dynamic cell by using a constant-temperature molecular dynamics simulation method. RxMC forward and reverse reaction steps are performed in the control cell only, while molecular dynamics steps are performed in both the dynamic cell and the control cell. The control cell, which acts as a sink and source reservoir, is maintained at reaction equilibrium conditions via the RxMC algorithm. The reaction ensemble molecular dynamics method is analogous to the grand canonical ensemble molecular dynamics technique, while using some elements of the osmotic molecular dynamics method, and so simulates conditions that directly relate to real, open systems. The accuracy and stability of the method is assessed by considering the ammonia synthesis reaction $N_2 + 3H_2 \rightleftharpoons 2NH_3$. It is shown to be a viable method for predicting the effects of nonideal environments on the dynamic properties (particularly diffusion) as well as reaction equilibria for chemically reacting mixtures.</p>					
15. SUBJECT TERMS molecular simulation, Monte Carlo, reaction ensemble					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UL	18. NUMBER OF PAGES 10	19a. NAME OF RESPONSIBLE PERSON John K. Brennan
a. REPORT UNCLASSIFIED	b. ABSTRACT UNCLASSIFIED	c. THIS PAGE UNCLASSIFIED			19b. TELEPHONE NUMBER (Include area code) 410-306-0678

Reaction ensemble molecular dynamics: Direct simulation of the dynamic equilibrium properties of chemically reacting mixtures

John K. Brennan,^{1,*} Martin Lísal,^{2,3} Keith E. Gubbins,⁴ and Betsy M. Rice¹

¹*U.S. Army Research Laboratory, Weapons and Materials Research Directorate, Aberdeen Proving Ground, Maryland 21005-5066, USA*

²*E. Hála Laboratory of Thermodynamics, Institute of Chemical Process Fundamentals, Academy of Sciences of the Czech Republic, 165 02 Prague 6, Czech Republic*

³*Department of Physics, J. E. Purkyne University, 400 96 Usti n. Lab., Czech Republic*

⁴*North Carolina State University, Department of Chemical Engineering, Raleigh, North Carolina 27695-7905, USA*
(Received 19 April 2004; published 15 December 2004)

A molecular simulation method to study the dynamics of chemically reacting mixtures is presented. The method uses a combination of stochastic and dynamic simulation steps, allowing for the simulation of both thermodynamic and transport properties. The method couples a molecular dynamics simulation cell (termed dynamic cell) to a reaction mixture simulation cell (termed control cell) that is formulated upon the reaction ensemble Monte Carlo (RxMC) method, hence the term reaction ensemble molecular dynamics. Thermodynamic and transport properties are calculated in the dynamic cell by using a constant-temperature molecular dynamics simulation method. RxMC forward and reverse reaction steps are performed in the control cell only, while molecular dynamics steps are performed in both the dynamic cell and the control cell. The control cell, which acts as a sink and source reservoir, is maintained at reaction equilibrium conditions via the RxMC algorithm. The reaction ensemble molecular dynamics method is analogous to the grand canonical ensemble molecular dynamics technique, while using some elements of the osmotic molecular dynamics method, and so simulates conditions that directly relate to real, open systems. The accuracy and stability of the method is assessed by considering the ammonia synthesis reaction $N_2 + 3H_2 \rightleftharpoons 2NH_3$. It is shown to be a viable method for predicting the effects of nonideal environments on the dynamic properties (particularly diffusion) as well as reaction equilibria for chemically reacting mixtures.

DOI: 10.1103/PhysRevE.70.061103

PACS number(s): 82.60.-s, 02.70.Ns, 05.10.Ln, 82.33.-z

I. INTRODUCTION

Predicting and understanding the physical effects of non-ideal conditions (strong intermolecular forces, nanostructured media, etc.) on chemical reaction equilibria is critical in many fields of science including mixture separation and purification using porous solids, catalysis, plasma physics, and shock physics. The reaction ensemble Monte Carlo method (RxMC) [1–3] is a powerful simulation tool for studying reaction mixtures and is uniquely capable of predicting shifts of reaction equilibria caused by such highly nonideal environments. The RxMC method requires as input only the intermolecular potentials and the ideal-gas partition functions for the reaction species that are present. Furthermore, the method does not require a reactive-type potential that mimics bond breakage and formation. Applications of the RxMC simulation method include reactive systems confined in porous materials [4–10], reactions of plasmas [11,12], reactions in supercritical fluid solvents [10], reactions under shock [13], and still others [14–17]. Deviations from the ideal-gas phase reaction equilibria caused by non-ideal conditions can be determined for quantities such as the fluid density, pressure, and species concentrations.

However, dynamic properties such as diffusion coefficients cannot be determined using the RxMC method. In this

work, we introduce a simulation method that can determine the dynamic properties of reaction mixtures at equilibrium. The method is akin to the grand canonical molecular dynamics method [18–20] and the osmotic molecular dynamics method [21,22]. These methods use a control cell to maintain the desired chemical potential while a dynamic cell is used to determine the dynamic properties. For the method introduced here, termed the reaction ensemble molecular dynamics (RxMD) method, the control cell is used to maintain the system at the reaction equilibrium conditions [1–3]. RxMC forward and reverse reaction steps are performed in the control cell only, while constant-temperature molecular dynamics steps are performed in both the dynamic cell and the control cell. The dynamic cell is in direct contact with the control cell so that particles are able to move freely between the cells. Since the entire simulation box (consisting of the control cell and the dynamic cell) is in thermodynamic equilibrium due to the molecular dynamics steps, reaction equilibrium conditions are established in the dynamic cell as a consequence of the physical contact between the control and dynamic cell. In this scenario, fluid properties in the dynamic cell are unaffected by the stochastic reaction steps occurring in the control cell. Therefore, dynamic quantities of reaction mixtures such as the velocity autocorrelation functions and the diffusion coefficients can be accurately determined in the dynamic cell. These dynamic properties are more precisely *dynamic equilibrium properties* since they describe correlations at different times along an equilibrium trajectory [23].

*Corresponding author. FAX: 410.306.1909. Electronic address: jkbrennan@arl.army.mil




FIG. 1. Schematic of the reaction ensemble molecular dynamics method. The model reaction $2A \rightleftharpoons B$ is occurring. Molecular dynamics particle displacement steps (\curvearrowright) occur in all cells, while reaction ensemble Monte Carlo reaction steps ($\blacktriangleleft \cdots \blacktriangleright$) occur only in the control cell. The dashed lines denote the portion of the dynamic cell in which the dynamic properties are calculated.

It is important to note further that, analogous to the RxMC method, the RxMD method predicts the *physical* effects on reaction equilibria, as opposed to predicting *chemical* effects. The RxMD method does not provide reaction rate information; separate methods must be used for this purpose (for a review of these methods see Santiso and Gubbins [24]). Despite these limitations, the RxMD method can provide unique insight into the molecular-level dynamic behavior of a wide variety of reacting systems. In particular, it enables the simultaneous study of the influence of nonideality and nanostructure on both diffusion and reaction equilibria.

II. METHODOLOGY

The reaction ensemble molecular dynamics method provides insight into the dynamic phenomena of reaction mixtures by combining a stochastic simulation method (RxMC) with a deterministic method (constant- T MD). The RxMC method can be performed at constant-volume or at constant-pressure conditions. The constant-volume version requires two types of Monte Carlo moves: (1) particle displacements and (2) forward and reverse reaction steps. Multiple reactions can be simulated simultaneously by including the forward and reverse reaction steps for each reaction in order to maintain stoichiometry. The constant-pressure version of the RxMC method requires the additional step of fluctuating the simulation cell volume to achieve the desired pressure. Further details of the RxMC method can be found elsewhere [1–3]. In this work, we demonstrate the RxMD method at constant-volume conditions; extension to constant-pressure conditions is straightforward.

The reaction ensemble molecular dynamics method is a direct extension of the reaction ensemble Monte Carlo method. In the RxMD method, Monte Carlo displacements steps are replaced by molecular dynamics time steps, while forward and reverse reaction steps are still performed in a Monte Carlo fashion. Reaction steps in the RxMC method require particle insertion, particle deletion, and/or particle identity exchange in the simulation cell. Such conditions are typically not suitable for the molecular dynamics technique since the deterministic pathway of the particle trajectories will be disrupted by such events. To avoid these adverse effects to the particle trajectories, the control cell and dynamic cell simulation setup described above is implemented. In practice, this entails positioning one-half of the control cell on each side of the dynamic cell. Such an arrangement allows for the use of periodic boundary conditions in all

directions. Moreover, total momentum is inherently conserved due to symmetry of the total simulation box. A schematic of the RxMD method is given in Fig. 1, where the model reaction $2A \rightleftharpoons B$ is occurring. In Fig. 1, molecular dynamics time steps are performed in both the control and dynamic cells while reaction steps are performed in the control cell only. Since the control cell is in direct contact with the dynamic cell, the particles are able to move freely between both cells. To minimize interface effects, the properties of the fluid in the dynamic cell are determined from an interior portion of the cell, which is away from the control-cell–dynamic-cell interface; i.e., at any time step only particles that reside within the interior portion are included in the properties calculation. The dashed lines in Fig. 1 are intended to illustrate this procedure. Last, particle velocities for newly inserted particles are drawn from a Maxwell-Boltzmann distribution at the appropriate temperature [23,25].

III. VERIFICATION: APPLICATION TO AMMONIA SYNTHESIS

We demonstrate the RxMD method using the ammonia synthesis reaction $N_2 + 3H_2 \rightleftharpoons 2NH_3$ at constant volume. Due to the illustrative nature of this work, the reacting species are modeled as simple spherical particles interacting through an exponential-6 potential, where electrostatics contributions are ignored [26]. The potential was truncated and shifted at 1.05 nm. The potential parameters are given in Table I, where the Lorentz-Berthelot combining rules [27] were used for unlike-pair interactions. Molecular partition functions were determined using JANAF thermochemical data tables [28]. All simulations were initiated from an initial mixture of 600 H_2 molecules and 200 N_2 molecules. For the state conditions simulated here, these relatively large system sizes result in large simulation cell volumes that help to minimize any interface effects caused by the dynamic-cell–control-cell boundary. The interior two-thirds of the dynamic cell were

TABLE I. Exponential-6 potential parameters [26].

Species	r_{core} (Å)	r_m (Å)	ε/k_B (K)	α
NH_3	1.12	3.72	244.9	12.0
N_2	1.03	4.17	97.1	13.0
H_2	1.25	3.49	30.4	11.2

TABLE II. Comparison of properties calculated by the reaction ensemble molecular dynamics method (RxMD) with the reaction ensemble Monte Carlo (RxMC) and molecular dynamics (MD) methods at various temperatures.^a

Method	U_{conf} (J/mol)	P [MPa]	density [g/cm ³]	$x(\text{NH}_3)^b$	$x(\text{N}_2)$	$x(\text{H}_2)$	$D(\text{NH}_3)$ (10 ⁸ m ² /s)
$T=300.0$ K							
RxMC	-3.9 ₂	0.0297 ₆	0.000191 ₆	0.882 ₃	0.0294 ₁	0.0882 ₁	NA ^c
RxMD dynamic cell	-3.5 ₆	0.0303 ₄	0.000202 ₈	0.885 ₄	0.0291 ₉	0.0852 ₆	552.1 ₄
control cell	-4.1 ₄	0.0311 ₈	0.000186 ₉	0.883 ₇	0.0301 ₅	0.0867 ₂	NA
MD-NVT	-3.8 ₄	0.0304 ₄	0.000191	0.882	0.0294	0.0882	552.8 ₅
$T=600.0$ K							
RxMC	-383.7 ₁₉	14.83 ₄	0.0382 ₇	0.526 ₁	0.118 ₄	0.355 ₂	NA
RxMD dynamic cell	-384.2 ₁₁	14.77 ₅	0.0399 ₈	0.527 ₃	0.115 ₈	0.357 ₄	3.584 ₆
control cell	-381.9 ₂₆	14.86 ₆	0.0387 ₉	0.526 ₁	0.116 ₃	0.358 ₅	NA
MD-NVT	-382.6 ₁₃	14.83 ₅	0.0382	0.526	0.118	0.355	3.578 ₈
$T=900.0$ K							
RxMC	-203.3 ₃₈	55.9 ₈	0.0668 ₇	0.178 ₁	0.205 ₂	0.616 ₃	NA
RxMD dynamic cell	-201.6 ₂₇	55.4 ₅	0.0654 ₈	0.181 ₅	0.197 ₄	0.622 ₈	2.781 ₉
control cell	-202.1 ₁₅	56.5 ₆	0.0660 ₆	0.173 ₃	0.209 ₇	0.619 ₂	NA
MD-NVT	-202.7 ₁₈	55.7 ₄	0.0668	0.178	0.205	0.616	2.788 ₇

^aReported uncertainties determined from block averages where, for example -383.7₁₉ implies -383.7 ± 1.9 [23].^b $x(i)$: mole fraction of species $i = N_i/N_{\text{total}}$, where N is the number of molecules.^cNA=not applicable to method.

used to calculate the fluid properties. Cubic simulation cells were used and periodic boundary conditions were imposed in all three directions for the total system. All reported pressures were calculated using the virial expression [25].

A standard *NVT* molecular dynamics method was employed with the equations of motion solved using the Verlet leapfrog algorithm [25] and implementing the damped force method of Brown and Clark to maintain constant temperature [29]. A time step of $\Delta t = 3.0$ fs was used for a total simulation time of 2 ns following an equilibration period. The ratio of attempted reaction steps per molecular dynamics steps was 40:1; this choice was based on a series of short simulation runs that were made during the development of the method, but may vary for different systems.

The ammonia synthesis reaction was simulated at a series of temperatures under known gas-phase conditions [4]. Comparisons of the quantities calculated by the RxMD method are made with two different methods. First, the thermodynamic quantities calculated from the RxMD approach (e.g., density, configurational energy, pressure, and species concentrations) are compared to quantities calculated by the RxMC approach. Second, the dynamic quantities of the RxMD method (e.g., velocity autocorrelation functions) are compared to quantities determined from an *NVT* molecular dynamics mixture simulation.

For a comparison of the thermodynamic properties, a constant-volume RxMC simulation was performed at the same temperature and initial mixture as the RxMD simulation. Table II presents a comparison of results for the RxMD and RxMC approaches, where quantities calculated in both the dynamic cell and the control cell are given. As expected, quantities in the dynamic cell and control cell are the same

within statistical uncertainty, ensuring that the RxMD has equilibrated properly. The quantities calculated using the RxMD and the RxMC approaches also agree within statistical uncertainty for all the temperatures considered.

For a comparison of the dynamic properties, *NVT* molecular dynamics simulations were performed at the temperature, volume, and species concentrations that were used or determined in the RxMC simulations. Self-diffusion coefficients (D) for each species were determined from the velocity autocorrelation functions [23] in both the *NVT*-MD simulations and in the dynamic cell of the RxMD simulations. Comparisons of the self-diffusion coefficients determined by both methods for NH_3 are given in Table II. Excellent agreement is evident between the two approaches, with similar agreement found for the self-diffusion coefficients of N_2 and H_2 .

IV. DISCUSSION

We have presented a simulation tool, the reaction ensemble molecular dynamics method, to study the dynamics of chemically reacting systems. The RxMD method is capable of predicting the physical effects of nonideal environments on dynamic properties of the equilibrium mixture. The RxMD method combines the reaction ensemble Monte Carlo method with the *NVT* molecular dynamics technique, resulting in a simulation method that mimics real, open systems for many experimental situations. The method as presented here effectively combines two simulation runs into a single simulation, since the same result could be achieved by running an RxMC simulation followed by a mixture *NVT*-MD simulation at the conditions determined from the RxMC result. The value of the RxMD method lies in its potential

application to other scenarios—e.g., where the dynamic cell is modeled as a porous solid and two control cells at different thermodynamic conditions are used. Such an arrangement mimics combined reaction and adsorption phenomena relevant to membrane reactors and fuel cells, as well as to various possible nanochemical devices [30].

ACKNOWLEDGMENTS

The authors gratefully acknowledge invaluable discussions with C. Heath Turner (University of Alabama). This

work was performed while J.K.B. received additional support from the U.S. Army Research Laboratory. M.L. acknowledges support from the Grant Agency of the Czech Republic (Grant No. 203/03/1588), and K.E.G. thanks the National Science Foundation (Grant No. CTS-0211792) for partial support of this work. Portions of the calculations reported in this work were performed at the Army Research Laboratory (Aberdeen Proving Ground, MD) and Engineer Research and Development Center (Vicksburg, MS) Major Shared Resource Centers.

-
- [1] J. K. Johnson, A. Z. Panagiotopoulos, and K. E. Gubbins, *Mol. Phys.* **81**, 717 (1994).
 - [2] W. R. Smith and B. Tříska, *J. Chem. Phys.* **100**, 3019 (1994).
 - [3] J. K. Johnson, *Adv. Chem. Phys.* **105**, 461 (1999).
 - [4] C. H. Turner, J. K. Johnson, and K. E. Gubbins, *J. Chem. Phys.* **114**, 1851 (2001).
 - [5] C. H. Turner, J. K. Brennan, J. K. Johnson, and K. E. Gubbins, in *Adsorption Science and Technology*, edited by D. D. Do (World Scientific, Singapore, 2001), p. 244.
 - [6] C. H. Turner, J. Pikunic, and K. E. Gubbins, *Mol. Phys.* **99**, 1991 (2001).
 - [7] M. Borówko and R. Zagórski, *J. Chem. Phys.* **114**, 5397 (2001).
 - [8] C. H. Turner, J. K. Brennan, J. Pikunic, and K. E. Gubbins, *Appl. Surf. Sci.* **196**, 366 (2002).
 - [9] C. H. Turner, J. K. Brennan, J. K. Johnson, and K. E. Gubbins, *J. Chem. Phys.* **116**, 2138 (2002).
 - [10] C. H. Turner and K. E. Gubbins, *J. Chem. Phys.* **119**, 6057 (2003).
 - [11] M. Lísál, W. R. Smith, and I. Nezbeda, *J. Chem. Phys.* **113**, 4885 (2000).
 - [12] M. Lísál, W. R. Smith, M. Bures, V. Vacek, and J. Navrátil, *Mol. Phys.* **100**, 2487 (2002).
 - [13] J. K. Brennan and B. M. Rice, *Phys. Rev. E* **66**, 021105 (2002).
 - [14] M. Lísál, I. Nezbeda, and W. R. Smith, *J. Chem. Phys.* **110**, 8597 (1999).
 - [15] M. Lísál, W. R. Smith, and I. Nezbeda, *J. Phys. Chem. B* **103**, 10 496 (1999).
 - [16] M. Lísál, W. R. Smith, and I. Nezbeda, *AIChE J.* **46**, 866 (2000).
 - [17] M. Lísál, W. R. Smith, and I. Nezbeda, *Fluid Phase Equilib.* **181**, 127 (2001).
 - [18] M. M. Cielinski, M.S. thesis, University of Maine at Orono, 1985.
 - [19] M. Lupkowski and F. van Swol, *J. Chem. Phys.* **95**, 1995 (1991).
 - [20] A. Papadopoulou, E. D. Becker, M. Lupkowski, and F. van Swol, *J. Chem. Phys.* **98**, 4897 (1993).
 - [21] R. L. Rowley, T. D. Shupe, and M. W. Shuck, *Mol. Phys.* **82**, 841 (1994).
 - [22] R. L. Rowley, M. W. Shuck, and J. C. Perry, *Mol. Phys.* **86**, 125 (1995).
 - [23] D. Frenkel and B. Smit, *Understanding Molecular Simulation* (Academic Press, San Diego, 2002).
 - [24] E. Santiso and K. E. Gubbins, *Mol. Simul.* **30**, 689 (2004).
 - [25] M. P. Allen and D. J. Tildesley, *Computer Simulation of Liquids* (Oxford University Press, Oxford, 1987).
 - [26] L. E. Fried and W. M. Howard, *J. Chem. Phys.* **109**, 7338 (1998).
 - [27] T. M. Reed and K. E. Gubbins, *Applied Statistical Mechanics* (McGraw-Hill, New York, 1973).
 - [28] M. W. Chase, C. A. Davies, J. R. Downey, D. J. Frurip, R. A. McDonald, and A. N. Syverud, *JANAF Thermochemical Tables*, 3rd ed. [*J. Phys. Chem. Ref. Data* **14**, 1551 (1985); **14**, 1260 (1985); **14**, 1293 (1985)].
 - [29] D. Brown and J. H. R. Clark, *Mol. Phys.* **51**, 1243 (1984).
 - [30] M. Lísál, J. K. Brennan, W. R. Smith, and F. R. Siperstein, *J. Chem. Phys.* **121**, 4901 (2004).

NO. OF
COPIES ORGANIZATION

1 DEFENSE TECHNICAL
(PDF INFORMATION CTR
ONLY) DTIC OCA
8725 JOHN J KINGMAN RD
STE 0944
FORT BELVOIR VA 22060-6218

1 US ARMY RSRCH DEV &
ENGRG CMD
SYSTEMS OF SYSTEMS
INTEGRATION
AMSRD SS T
6000 6TH ST STE 100
FORT BELVOIR VA 22060-5608

1 DIRECTOR
US ARMY RESEARCH LAB
IMNE ALC IMS
2800 POWDER MILL RD
ADELPHI MD 20783-1197

3 DIRECTOR
US ARMY RESEARCH LAB
AMSRD ARL CI OK TL
2800 POWDER MILL RD
ADELPHI MD 20783-1197

ABERDEEN PROVING GROUND

1 DIR USARL
AMSRD ARL CI OK TP (BLDG 4600)

NO. OF
COPIES ORGANIZATION

ABERDEEN PROVING GROUND

5 DIR USARL
 AMSRD ARL WM BD
 J BRENNAN